



I'm not robot



**Continue**

## Easl hepatitis b guidelines

Hepatitis B is caused by a virus. It is very contagious and heartbreaking. Hepatitis A and B are very similar when it comes to symptoms. They are both acute and can last up to six months. However, hepatitis B can also be chronic. The incubation period for hepatitis B is also longer. Although the infection can spread in the same way, the causes are different. Continue reading to learn more about hepatitis B along with its causes and treatments. If you have been infected with the hepatitis B virus, you can experience symptoms anywhere from one to four months later. Even after the infection clears, you can still exhibit symptoms for up to two weeks. Symptoms of hepatitis B can range from moderate to severe and may include fever, weakness, fatigue, and loss of appetite along with nausea and vomiting. Another common sign of hepatitis B is joint pain or abdominal pain, especially in the upper right side under the ribs where the liver is located. Jaundice, which is characterized with skin yellow and white your eyes is another symptom of liver infection. However, some people, mostly young children, may experience any symptoms at all. You may already know you are exposed to infectious viruses, in which case you should contact the doctor immediately. You can take preventive treatment within the first 24 hours of a relationship that may lower the chance of infection. If you are unsure of your condition, but have any signs or symptoms that have been mentioned before, please visit your doctor as soon as possible. The virus known as HBV causes hepatitis B. It is highly contagious, but does not spread through coughing or sneezing. On the other hand, it is spread through blood, semen, or body fluids. Therefore, sexual intercourse with someone with a virus either through saliva, semen, or other secretaries, may allow the virus to enter your body. A pregnant woman infected with HBV can also pass the virus to her baby during childbirth. Other causes of hepatitis B involve the sharing of needles. Using contaminated needles can spread infection either through the use of drugs or instant needles. Healthcare workers and others around the blood have a higher risk of contracting the infection. It is important to always use a clean needle and never share it with others. There are two types of HBV. Acute is a short-term version of the infection that lasts less than six months. On the other hand, a long-heated

